

GENERAL HEART CONSTRUCTION ON A TRIANGULATED CATEGORY (I): UNIFYING t -STRUCTURES AND CLUSTER TILTING SUBCATEGORIES

HIROYUKI NAKAOKA

ABSTRACT. In the paper of Keller and Reiten, it was shown that the quotient of a triangulated category (with some conditions) by a cluster tilting subcategory becomes an abelian category. After that, Koenig and Zhu showed in detail, how the abelian structure is given on this quotient category, in a more abstract setting. On the other hand, as is well known since 1980s, the heart of any t -structure is abelian. We unify these two construction by using the notion of a cotorsion pair. To any cotorsion pair in a triangulated category, we can naturally associate an abelian category, which gives back each of the above two abelian categories, when the cotorsion pair comes from a cluster tilting subcategory, or is a t -structure, respectively.

1. INTRODUCTION

Throughout this paper, we fix a triangulated category $\mathcal{C}$.

For any category $\mathcal{K}$, we write abbreviately $K \in \mathcal{K}$, to indicate that K is an object of $\mathcal{K}$.

For any $K, L \in \mathcal{K}$, let $\mathcal{K}(K, L)$ denote the set of morphisms from K to L .

If $\mathcal{M}, \mathcal{N}$ are full subcategories of $\mathcal{K}$, then $\mathcal{K}(\mathcal{M}, \mathcal{N}) = 0$ means that $\mathcal{K}(M, N) = 0$ for any $M \in \mathcal{M}$ and $N \in \mathcal{N}$.

Similarly, $\mathcal{K}(K, \mathcal{N}) = 0$ means $\mathcal{K}(K, N) = 0$ for any $N \in \mathcal{N}$.

When $\mathcal{K}$ admits Ext^ℓ , similarly for $\text{Ext}^\ell(\mathcal{M}, \mathcal{N}) = 0$ and $\text{Ext}^\ell(K, \mathcal{N}) = 0$ for any integer ℓ .

As is well known, if $(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$ is a t -structure on $\mathcal{C}$, then its heart $\mathcal{H} = \mathcal{T}^{\leq 0} \cap \mathcal{T}^{\geq 0}$ becomes an abelian category [BBD]. Put $\mathcal{T}^{\leq n} := \mathcal{T}^{\leq 0}[-n]$, $\mathcal{T}^{\geq n} := \mathcal{T}^{\geq 0}[-n]$ for an integer n . By definition, a t -structure is a pair of full additive thick subcategories $(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$ of $\mathcal{C}$ satisfying

- (t-1) $\mathcal{C}(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 1}) = 0$,
- (t-2) $\mathcal{T}^{\leq 0} \subseteq \mathcal{T}^{\leq 1}$ and $\mathcal{T}^{\geq 1} \subseteq \mathcal{T}^{\geq 0}$,
- (t-3) For any $C \in \mathcal{C}$, there exists a distinguished triangle

$$X \rightarrow C \rightarrow Y \rightarrow X[1] \quad (X \in \mathcal{T}^{\leq 0}, Y \in \mathcal{T}^{\geq 1}).$$

On the other hand, in [KZ], Koenig and Zhu showed that for any cluster tilting subcategory $\mathcal{T}$ of $\mathcal{C}$, the quotient category $\mathcal{C}/\mathcal{T}$ carries a naturally induced abelian

The author wishes to thank Professor Toshiyuki Katsura for his encouragement.

The author wishes to thank his colleague Professor Noriyuki Abe. This work was never possible without his advices.

The author wishes to thank Professor Bernhard Keller and Professor Osamu Iyama for their useful comments and advices, especially on the terminology.

structure. Originally, an equivalence between the quotient of a triangulated category and a certain module category was shown in [BMR] for the quotient by a full additive thick subcategory associated to a tilting object in $\mathcal{C}$, and then in [KR], for the quotients by a cluster tilting subcategory.

By definition ([KR], [KZ]), a full additive thick subcategory $\mathcal{T}$ of $\mathcal{C}$ is a cluster tilting subcategory if it satisfies

- (s-1) $\mathcal{T}$ is functorially finite (cf. [BR]),
- (s-2) $C \in \mathcal{T}$ if and only if $\text{Ext}^1(C, \mathcal{T}) = 0$,
- (s-3) $C \in \mathcal{T}$ if and only if $\text{Ext}^1(\mathcal{T}, C) = 0$.

By (s-1), it can be easily shown that a cluster tilting subcategory also satisfies the following for any $C \in \mathcal{C}$:

(s-1)' For any $C \in \mathcal{C}$, there exists a distinguished triangle

$$T \rightarrow C \rightarrow T'[1] \rightarrow T[1] \quad (T, T' \in \mathcal{T}).$$

To unify these two, we introduce the notion of a *cotorsion pair* (Definition 2.1). To any cotorsion pair $(\mathcal{U}, \mathcal{V})$ in $\mathcal{C}$, we can naturally associate a subfactor category $\underline{\mathcal{H}} := (\underline{\mathcal{C}}^+ \cap \underline{\mathcal{C}}^-) / (\mathcal{U} \cap \mathcal{V})$. As a main theorem, we show $\underline{\mathcal{H}}$ carries an induced abelian structure (Theorem 6.4). In fact, this construction generalizes the above abelian categories, in the following sense (Proposition 2.6 and Example 3.8):

- If $(\mathcal{U}, \mathcal{V})$ satisfies $\mathcal{V} \subseteq \mathcal{V}[1]$, then $(\mathcal{U}[-1], \mathcal{V}[1])$ becomes a t -structure, and $\underline{\mathcal{H}}$ agrees with the heart of $(\mathcal{U}[-1], \mathcal{V})$.
- If $(\mathcal{U}, \mathcal{V})$ satisfies $\mathcal{U} = \mathcal{V}$, then $\mathcal{T} := \mathcal{U} = \mathcal{V}$ becomes a cluster tilting subcategory of $\mathcal{C}$, and $\underline{\mathcal{H}}$ agrees with $\mathcal{C}/\mathcal{T}$.

2. PRELIMINARIES

Definition 2.1. Let $\mathcal{U}$ and $\mathcal{V}$ be full additive thick subcategories of $\mathcal{C}$. We call $(\mathcal{U}, \mathcal{V})$ a *cotorsion pair* if it satisfies the following.

- (1) $\text{Ext}^1(\mathcal{U}, \mathcal{V}) = 0$,
- (2) For any $C \in \mathcal{C}$, there exists a (not necessarily unique) distinguished triangle

$$U \rightarrow C \rightarrow V[1] \rightarrow U[1]$$

satisfying $U \in \mathcal{U}$ and $V \in \mathcal{V}$.

For any cotorsion pair $(\mathcal{U}, \mathcal{V})$, put

$$\mathcal{W} := \mathcal{U} \cap \mathcal{V}.$$

Remark 2.2. A pair $(\mathcal{U}, \mathcal{V})$ of full additive thick subcategories of $\mathcal{C}$ is a cotorsion pair if and only if $(\mathcal{U}[-1], \mathcal{V})$ is a *torsion theory* (or a *torsion pair*) in [IY]. (Unlike [BR], it does not require the shift-closedness.) In this sense, a cotorsion pair is nothing other than a torsion theory with $\mathcal{U}$ shifted by -1 , and thus not a new notion. However we prefer the above definition, just for the sake of the duality in the index of shifts.

Remark 2.3. A pair $(\mathcal{U}, \mathcal{V})$ of full additive subcategories of $\mathcal{C}$ is a cotorsion pair if and only if it satisfies the following conditions for any $C \in \mathcal{C}$.

- (1) C belongs to $\mathcal{U}$ if and only if $\text{Ext}^1(C, \mathcal{V}) = 0$,
- (2) C belongs to $\mathcal{V}$ if and only if $\text{Ext}^1(\mathcal{U}, C) = 0$,
- (3) For any $C \in \mathcal{C}$, there exists a (not necessarily unique) distinguished triangle

$$U \rightarrow C \rightarrow V[1] \rightarrow U[1]$$

satisfying $U \in \mathcal{U}$ and $V \in \mathcal{V}$.

Remark 2.4. Let $(\mathcal{U}, \mathcal{V})$ be a cotorsion pair in $\mathcal{C}$.

- (1) For any $C \in \mathcal{C}$ and any $n \in \mathbb{Z}$, C admits a distinguished triangle

$$U[n] \rightarrow C \rightarrow V[n+1] \rightarrow U[n+1]$$

with $U \in \mathcal{U}, V \in \mathcal{V}$.

- (2) For any $n \in \mathbb{Z}$, each $\mathcal{U}[n]$ and $\mathcal{V}[n]$ is closed under extensions and direct summands.
 (3) $\mathcal{W}$ becomes an additive full subcategory of $\mathcal{C}$, closed under direct sums and direct summands, satisfying

$$\text{Ext}^1(\mathcal{W}, \mathcal{W}) = 0.$$

Example 2.5.

- (1) If $(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$ is a t -structure on $\mathcal{C}$, then

$$(\mathcal{U}, \mathcal{V}) := (\mathcal{T}^{\leq -1}, \mathcal{T}^{\geq 1})$$

becomes a cotorsion pair. In this case, $\mathcal{W} = 0$.

- (2) If $\mathcal{T}$ is a cluster tilting subcategory of $\mathcal{C}$, then

$$(\mathcal{U}, \mathcal{V}) := (\mathcal{T}, \mathcal{T})$$

becomes a cotorsion pair. In this case, $\mathcal{W} = \mathcal{T}$.

Remark that $\text{Ext}^1(\mathcal{U}, \mathcal{V}) = 0$ implies $\mathcal{U} \cap \mathcal{V}[1] = 0$. The following proposition shows that t -structures and cluster tilting subcategories are characterized as two extremal examples of cotorsion pairs.

Proposition 2.6. *Let $(\mathcal{U}, \mathcal{V})$ be a cotorsion pair.*

- (1) $\mathcal{V} \subseteq \mathcal{V}[1]$ if and only if $(\mathcal{U}, \mathcal{V}) = (\mathcal{T}^{\leq -1}, \mathcal{T}^{\geq 1})$ for a t -structure $(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$.
 (2) $\mathcal{V} = \mathcal{U}$ if and only if $(\mathcal{U}, \mathcal{V}) = (\mathcal{T}, \mathcal{T})$ for a cluster tilting subcategory $\mathcal{T}$.

Proof. (1) This immediately follows from the definition of a t -structure.

(2) The ‘only if’ part is trivial. To show the converse, put $\mathcal{T} := \mathcal{U} = \mathcal{V}$. As in [KZ] (Definition 3.1 and Lemma 3.2), it suffices to show $\mathcal{T}$ is contravariantly finite. But this immediately follows from the fact that any object $C \in \mathcal{C}$ admits a distinguished triangle

$$T \rightarrow C \rightarrow T'[1] \rightarrow T[1]$$

for some $T \in \mathcal{U} = \mathcal{T}$ and $T' \in \mathcal{V} = \mathcal{T}$. □

Definition 2.7. (cf. §§2.1 in [KZ]) Let $(\mathcal{U}, \mathcal{V})$ be a cotorsion pair in $\mathcal{C}$, and $\mathcal{W} := \mathcal{U} \cap \mathcal{V}$. We denote the quotient of $\mathcal{C}$ by $\mathcal{W}$ as

$$\underline{\mathcal{C}} := \mathcal{C}/\mathcal{W}.$$

Namely, $\text{Ob}(\underline{\mathcal{C}}) = \text{Ob}(\mathcal{C})$, and for any $A, B \in \mathcal{C}$,

$$\underline{\mathcal{C}}(A, B) := \mathcal{C}(A, B)/\{f \in \mathcal{C}(A, B) \mid f \text{ factors through some } W \in \mathcal{W}\}.$$

For any morphism $f \in \mathcal{C}(A, B)$, we denote its image in $\underline{\mathcal{C}}(A, B)$ by $\underline{f}$. This defines an additive functor

$$(\underline{\quad}): \mathcal{C} \rightarrow \underline{\mathcal{C}}.$$

Since $\mathcal{U} \supseteq \mathcal{W}$ and $\mathcal{V} \supseteq \mathcal{W}$, we also have additive full subcategories of $\underline{\mathcal{C}}$

$$\underline{\mathcal{U}} := \mathcal{U}/\mathcal{W} \quad \text{and} \quad \underline{\mathcal{V}} := \mathcal{V}/\mathcal{W}.$$

Remark 2.8. Since $\mathcal{W}$ is closed under direct summands, for any $C \in \mathcal{C}$ we have

$$C \cong 0 \text{ in } \underline{\mathcal{C}} \iff C \in \mathcal{W}.$$

Since $\text{Ext}^1(\mathcal{W}, \mathcal{W}) = 0$, Theorem 2.3 in [KZ] can be applied. Compare with Proposition 6.1.

Fact 2.9. (Theorem 2.3 in [KZ]) Let $\mathcal{C}$, $(\mathcal{U}, \mathcal{V})$ and $\mathcal{W}$ be as above. For any distinguished triangle

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} X[1],$$

the following hold.

- (1) $\underline{g}$ is epimorphic in $\underline{\mathcal{C}}$ if and only if $\underline{h} = 0$,
- (2) $\underline{g}$ is monomorphic in $\underline{\mathcal{C}}$ if and only if $\underline{f} = 0$.

Let $(\mathcal{U}, \mathcal{V})$ be a cotorsion pair in the following.

Proposition 2.10.

- (1) For any $U \in \mathcal{U}$, $C \in \mathcal{C}$ and any $f \in \mathcal{C}(U[-1], C)$, if $\underline{f} = 0$, then $f = 0$.
Namely, we have

$$\underline{\mathcal{C}}(U[-1], C) = \mathcal{C}(U[-1], C) = \text{Ext}^1(U, C).$$

In particular, $C \in \mathcal{V}$ if and only if $\underline{\mathcal{C}}(U[-1], C) = 0$ for any $U \in \mathcal{U}$.

- (2) Dually, for any $V \in \mathcal{V}$ and any $C \in \mathcal{C}$, we have

$$\underline{\mathcal{C}}(C, V[1]) = \mathcal{C}(C, V[1]) = \text{Ext}^1(C, V).$$

In particular, $C \in \mathcal{U}$ if and only if $\underline{\mathcal{C}}(C, V[1]) = 0$ for any $V \in \mathcal{V}$.

Proof. (1) By definition, $\underline{f} = 0$ if and only if f factors through some $W \in \mathcal{W}$. Since $\mathcal{C}(U[-1], W) \subseteq \text{Ext}^1(\mathcal{U}, \mathcal{V}) = 0$, this implies $f = 0$.

$$\begin{array}{ccc} & W & \\ \nearrow 0 & \circlearrowleft & \searrow \\ U[-1] & \xrightarrow{f} & C \end{array}$$

(2) is shown dually. □

Lemma 2.11. For any cotorsion pair $(\mathcal{U}, \mathcal{V})$, we have

$$\underline{\mathcal{C}}(\mathcal{U}, \mathcal{V}) = 0.$$

Proof. Let $f \in \mathcal{C}(U, V)$ be a morphism, where $U \in \mathcal{U}$ and $V \in \mathcal{V}$. By condition (2) in Definition 2.1, we can form a distinguished triangle

$$V' \rightarrow U' \xrightarrow{u} V \xrightarrow{v} V'[1]$$

where $U' \in \mathcal{U}$ and $V' \in \mathcal{V}$. Since $\mathcal{V}$ is extension-closed (Remark 2.4), U' satisfies $U' \in \mathcal{U} \cap \mathcal{V} = \mathcal{W}$. Since $v \circ f = 0$ by $\text{Ext}^1(U, V') = 0$, f factors through U' .

$$\begin{array}{ccccc} & & U & & \\ & \swarrow & \downarrow f & \searrow & \\ V' & \longrightarrow & U' & \longrightarrow & V \xrightarrow{v} V'[1] \end{array}$$

Since $U' \in \mathcal{W}$, this means $\underline{f} = 0$. □

3. DEFINITION OF $\mathcal{C}^+$ AND $\mathcal{C}^-$

Lemma 3.1. *Let $f: A \rightarrow B$ be any morphism in $\mathcal{C}$.*

(1) *Let*

$$\begin{aligned} U_A &\xrightarrow{u_A} A \rightarrow V_A[1] \rightarrow U_A[1] \\ U_B &\xrightarrow{u_B} B \rightarrow V_B[1] \rightarrow U_B[1] \end{aligned}$$

be any distinguished triangles satisfying $U_A, U_B \in \mathcal{U}$ and $V_A, V_B \in \mathcal{V}$. Then there exists a morphism $f_U \in \mathcal{C}(U_A, U_B)$ such that

$$f \circ u_A = u_B \circ f_U.$$

$$\begin{array}{ccc} U_A & \xrightarrow{u_A} & A \\ f_U \downarrow & \circlearrowleft & \downarrow f \\ U_B & \xrightarrow{u_B} & B \end{array}$$

Moreover, f_U with this property is unique in $\mathcal{C}(U_A, U_B)$.

(2) *Dually, for any distinguished triangles*

$$\begin{aligned} U'_A[-1] &\rightarrow A \rightarrow V'_A \rightarrow U'_A \\ U'_B[-1] &\rightarrow B \rightarrow V'_B \rightarrow U'_B \end{aligned}$$

with $U'_A, U'_B \in \mathcal{U}$ and $V'_A, V'_B \in \mathcal{V}$, there exists a morphism $f'_V \in \mathcal{C}(V'_A, V'_B)$ compatible with f , uniquely up to $\mathcal{W}$.

Proof. We only show (1). Existence immediately follows from $\mathcal{C}(U_A, V_B[1]) = 0$. Moreover if f_U^1 and f_U^2 in $\mathcal{C}(U_A, U_B)$ satisfies

$$f_U^1 \circ u_A = u_B \circ f_U = f_U^2 \circ u_A,$$

then by $(f_U^1 - f_U^2) \circ u_A = 0$, there exists $w \in \mathcal{C}(U_A, V_B)$ such that $f_U^1 - f_U^2$ factors w .

$$\begin{array}{ccccccc} & & U_A & \xrightarrow{u_A} & A & & \\ & \swarrow w & \downarrow & & \downarrow f & & \\ V_B & \xrightarrow{\quad} & U_B & \xrightarrow{\quad} & B & \xrightarrow{\quad} & V_B[1] \end{array}$$

By Lemma 2.11 we have $\underline{w} = 0$, and thus $\underline{f_U^1} = \underline{f_U^2}$. □

Proposition 3.2. *Let C be any object in $\mathcal{C}$.*

(1) *For any distinguished triangles*

$$\begin{aligned} U &\xrightarrow{u} C \rightarrow V[1] \rightarrow U[1] \\ U' &\xrightarrow{u'} C \rightarrow V'[1] \rightarrow U'[1] \end{aligned}$$

satisfying $U, U' \in \mathcal{U}$ and $V, V' \in \mathcal{V}$, there exists a morphism $s \in \mathcal{C}(U, U')$ compatible with u and u' , such that $\underline{s}$ is an isomorphism.

$$\begin{array}{ccc} U & & U \\ \parallel \downarrow \underline{s} & & \downarrow s \\ U' & & U' \end{array} \quad \begin{array}{ccc} & u & \\ & \searrow & \\ & \circlearrowleft & \\ & \nearrow u' & \\ & C & \end{array}$$

(2) Dually, those V appearing in distinguished triangles

$$U[-1] \rightarrow C \rightarrow V \rightarrow U \quad (U \in \mathcal{U}, V \in \mathcal{V})$$

are isomorphic in $\underline{\mathcal{C}}$.

Proof. This immediately follows from Lemma 3.1. $\square$

Corollary 3.3. *For any $C \in \mathcal{C}$, the following are equivalent.*

(1) *There exists a distinguished triangle*

$$W_0 \rightarrow C \rightarrow V_0[1] \rightarrow W_0[1]$$

such that $W_0 \in \mathcal{W}, V_0 \in \mathcal{V}$.

(2) *Any distinguished triangle*

$$U \rightarrow C \rightarrow V[1] \rightarrow U[1] \quad (U \in \mathcal{U}, V \in \mathcal{V})$$

satisfies $U \in \mathcal{W}$.

Proof. Suppose (1) holds. By Proposition 3.2, we have $U \cong W_0$ in $\underline{\mathcal{C}}$. By Remark 2.8, this means $U \in \mathcal{W}$. The converse is trivial. $\square$

Dually, we have the following:

Corollary 3.4. *For any $C \in \mathcal{C}$, the following are equivalent.*

(1) *There exists a distinguished triangle*

$$U_0[-1] \rightarrow C \rightarrow W_0 \rightarrow U_0$$

such that $U_0 \in \mathcal{U}, W_0 \in \mathcal{W}$.

(2) *Any distinguished triangle*

$$U[-1] \rightarrow C \rightarrow V \rightarrow U \quad (U \in \mathcal{U}, V \in \mathcal{V})$$

satisfies $V \in \mathcal{W}$.

Definition 3.5.

- (1) $\mathcal{C}^+$ is defined to be the full subcategory of $\mathcal{C}$, consisting of objects satisfying equivalent conditions of Corollary 3.3.
- (2) $\mathcal{C}^-$ is defined to be the full subcategory of $\mathcal{C}$, consisting of objects satisfying equivalent conditions of Corollary 3.4.

Remark 3.6. The following are satisfied.

- (1) Each of $\mathcal{C}^+$ and $\mathcal{C}^-$ is an additive full subcategory of $\mathcal{C}$ containing $\mathcal{W}$.
- (2) $\mathcal{C}^+ \supseteq \mathcal{V}[1]$.
- (3) $\mathcal{C}^- \supseteq \mathcal{U}[-1]$.

Definition 3.7. For any cotorsion pair $(\mathcal{U}, \mathcal{V})$, put $\mathcal{H} := \mathcal{C}^+ \cap \mathcal{C}^-$. Since $\mathcal{H} \supseteq \mathcal{W}$, we have an additive full subcategory

$$\underline{\mathcal{H}} := \mathcal{H}/\mathcal{W} \subseteq \underline{\mathcal{C}},$$

which we call *the heart* of $(\mathcal{U}, \mathcal{V})$.

Example 3.8.

- (1) If $(\mathcal{U}, \mathcal{V}) = (\mathcal{T}^{\leq -1}, \mathcal{T}^{\geq 1})$, where $(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$ is a t -structure, then we have

$$\begin{aligned}\mathcal{C}^- &= \mathcal{T}^{\leq -1}[-1] = \mathcal{T}^{\leq 0}, \\ \mathcal{C}^+ &= \mathcal{T}^{\geq 1}[1] = \mathcal{T}^{\geq 0}, \\ \underline{\mathcal{H}} &= \mathcal{H} = \mathcal{T}^{\leq 0} \cap \mathcal{T}^{\geq 0}.\end{aligned}$$

Thus the definition of the heart agrees with that of a t -structure. Thus $\underline{\mathcal{H}}$ is abelian, and admits a cohomological functor $H^0: \mathcal{C} \rightarrow \underline{\mathcal{H}}$ (cf. [BBD]).

- (2) If $\mathcal{U} = \mathcal{V} = \mathcal{T}$ is a cluster tilting subcategory of $\mathcal{C}$, then we have

$$\begin{aligned}\mathcal{C}^+ &= \mathcal{C}^- = \mathcal{H} = \mathcal{C}, \\ \underline{\mathcal{H}} &= \mathcal{C}/\mathcal{T}.\end{aligned}$$

By [KZ], $\underline{\mathcal{H}}$ becomes an abelian category, and the quotient functor $(\underline{\quad}): \mathcal{C} \rightarrow \mathcal{C}/\mathcal{T} = \underline{\mathcal{H}}$ is cohomological.

4. EXISTENCE OF (CO)REFLECTIONS

Since $\mathcal{C}^+ \cap \mathcal{C}^- = \mathcal{H} \supseteq \mathcal{W}$, we have additive full subcategories of $\underline{\mathcal{C}}$

$$\underline{\mathcal{C}}^+ := \mathcal{C}^+/\mathcal{W} \quad \text{and} \quad \underline{\mathcal{C}}^- := \mathcal{C}^-/\mathcal{W}.$$

$$\begin{array}{ccccc} & & \underline{\mathcal{C}}^+ & & \\ & \nearrow & \hookrightarrow & \searrow & \\ \underline{\mathcal{H}} & & \circ & & \underline{\mathcal{C}} \\ & \searrow & \hookrightarrow & \nearrow & \\ & & \underline{\mathcal{C}}^- & & \end{array}$$

Definition 4.1. (Definition 3.1.1 in [B]) Let $\mathcal{A}$ and $\mathcal{B}$ be categories, and $F: \mathcal{A} \rightarrow \mathcal{B}$ be a functor. For any $B \in \mathcal{B}$, a reflection of B along F is a pair (R_B, η_B) of $R_B \in \mathcal{A}$ and $\eta_B \in \mathcal{B}(B, F(R_B))$, satisfying the following universality:

(*) For any $A \in \mathcal{A}$ and any $b \in \mathcal{B}(B, F(A))$, there exists a unique morphism $a \in \mathcal{A}(R_B, A)$ such that $F(a) \circ \eta_B = b$.

$$\begin{array}{ccc} B & \xrightarrow{\eta_B} & F(R_B) \\ & \searrow b \quad \swarrow F(a) & \\ & F(A) & \end{array}$$

A *coreflection* is defined dually.

Construction 4.2. For any $C \in \mathcal{C}$, consider a diagram

$$\begin{array}{ccccc} U'_C[-1] & & & & \\ \downarrow u'_C & \searrow w_C & & & \downarrow v_C \\ & \circ & & & V_C[1] \\ & & C & \xrightarrow{v_C} & \\ U_C & \xrightarrow{u_C} & & \searrow y_C & \\ & \circ & & & \\ & & Z_C & & \\ \downarrow v'_C & \nearrow x_C & & & \\ V'_C & & & & \end{array}$$

where

$$\begin{aligned} U_C &\xrightarrow{u_C} C \xrightarrow{v_C} V_C[1] \rightarrow U_C[1] \\ U'_C[-1] &\xrightarrow{u'_C} U_C \xrightarrow{v'_C} V'_C \rightarrow U'_C \\ U'_C[-1] &\xrightarrow{w_C} C \xrightarrow{z_C} Z_C \rightarrow U'_C \end{aligned}$$

are distinguished triangles, satisfying $U_C, U'_C \in \mathcal{U}$ and $V_C, V'_C \in \mathcal{V}$. Since $\mathcal{U}$ is extension-closed, we have $V'_C \in \mathcal{W}$. By the octahedron axiom,

$$V'_C \xrightarrow{x_C} Z_C \xrightarrow{y_C} V_C[1] \rightarrow V'_C$$

also becomes a distinguished triangle. Thus Z_C belongs to $\mathcal{C}^+$.

Proposition 4.3. *In the notation of Construction 4.2, for any $C \in \mathcal{C}$,*

$$\underline{z}_C: C \rightarrow Z_C$$

gives a reflection of C along $\underline{\mathcal{C}}^+ \hookrightarrow \underline{\mathcal{C}}$.

Proof. Let Y be any object in $\mathcal{C}^+$, and let $y \in \mathcal{C}(C, Y)$ be any morphism. It suffices to show that there exists a unique morphism

$$\underline{q} \in \underline{\mathcal{C}}(Z_C, Y)$$

such that $\underline{q} \circ \underline{z}_C = \underline{y}$.

$$\begin{array}{ccc} C & \xrightarrow{\underline{z}_C} & Z_C \\ & \searrow y & \swarrow \underline{q} \\ & Y & \end{array}$$

In fact, q can be chosen to satisfy $q \circ z_C = y$.

First, we show the existence. Since

$$U'_C[-1] \xrightarrow{w_C} C \xrightarrow{z_C} Z_C \rightarrow U'_C$$

is a distinguished triangle, it suffices to show $y \circ w_C = 0$.

Let

$$U_Y \xrightarrow{u_Y} Y \xrightarrow{v_Y} V_Y[1] \rightarrow U_Y[1]$$

be a distinguished triangle such that $U_Y \in \mathcal{W}, V_Y \in \mathcal{V}$. By $\mathcal{C}(U_C, V_Y[1]) = 0$, there exists a morphism $y_C \in \mathcal{C}(U_C, U_Y)$ such that $y \circ u_C = u_Y \circ y_U$.

$$\begin{array}{ccccc} & & U'_C[-1] & & \\ & \swarrow u'_C & & \searrow w_C & \\ U_C & & \circ & & C \\ & \swarrow u_C & & \searrow y & \\ & U_Y & & Y & \\ & \swarrow u_Y & & & \end{array}$$

Since $U_Y \in \mathcal{W}$, we have

$$\underline{y} \circ \underline{w}_C = \underline{u}_Y \circ \underline{y}_U \circ \underline{u}'_C = 0.$$

By Proposition 2.10, we obtain $y \circ w_C = 0$.

To show the uniqueness, suppose $q, q' \in \mathcal{C}(Z_C, Y)$ satisfies

$$\underline{q} \circ \underline{z}_C = \underline{y} = \underline{q}' \circ \underline{z}_C.$$

$$\begin{array}{ccc} C & \xrightarrow{z_C} & Z_C \\ & \searrow y \quad \circ \quad \swarrow q, q' & \\ U_Y & \xrightarrow{u_Y} Y & \xrightarrow{v_Y} V_Y[1] \end{array}$$

Since $\underline{v}_Y \circ (\underline{q} - \underline{q}') \circ \underline{z}_C = 0$ in $\underline{\mathcal{C}}(C, V_Y[1])$, it follows $v_Y \circ (q - q') \circ z_C = 0$ by Proposition 2.10.

Thus we have a morphism of triangles

$$\begin{array}{ccccc} C & \xrightarrow{z_C} & Z_C & \longrightarrow & U'_C \\ \downarrow & \circ & \downarrow q-q' & \circ & \downarrow \\ U_Y & \xrightarrow{u_Y} & Y & \xrightarrow{v_Y} & V_Y[1]. \end{array}$$

Since $\mathcal{C}(U'_C, V_Y[1]) = 0$, this implies $v_Y \circ (q - q') = 0$. Thus $q - q'$ factors through $U_Y \in \mathcal{W}$, which means $\underline{q} = \underline{q}'$. $\square$

Corollary 4.4. (*Proposition 3.1.2 and Proposition 3.1.3 in [B]*)

- (1) Since (Z_C, z_C) is a reflection of C along $\underline{\mathcal{C}}^+ \hookrightarrow \underline{\mathcal{C}}$, it is determined up to a canonical isomorphism in $\underline{\mathcal{C}}^+$.
- (2) As in [B], if we allow the axiom of choice, we obtain a left adjoint $\sigma: \underline{\mathcal{C}} \rightarrow \underline{\mathcal{C}}^+$ of the inclusion $\underline{\mathcal{C}}^+ \xrightarrow{i^+} \underline{\mathcal{C}}$. If we denote the adjunction by $\eta: \text{Id}_{\underline{\mathcal{C}}} \Rightarrow i^+ \circ \sigma$, then there exists a canonical isomorphism $Z_C \cong \sigma(C)$ in $\underline{\mathcal{C}}$, compatible with $\underline{z}_C$ and η_C .

$$\begin{array}{ccc} C & \xrightarrow{z_C} & Z_C \\ \eta_C \searrow & \circ & \swarrow \cong \\ & \sigma(C) & \end{array}$$

Dually, we have the following:

Remark 4.5. For any $C \in \mathcal{C}$, if we take a diagram

$$\begin{array}{ccccc} U''_C[-1] & \longrightarrow & K_C & \longrightarrow & U'''_C \\ & \searrow \circ & \downarrow k_C \quad \circ & & \swarrow \\ & & C & \longrightarrow & V''_C \\ & \searrow \circ & & \swarrow \circ & \\ & & V'''_C[1] & & \end{array} \quad (U''_C, U'''_C \in \mathcal{U}, V''_C, V'''_C \in \mathcal{V})$$

where all 180° composition of arrows are distinguished triangles, then $\underline{k}_C: K_C \rightarrow C$ gives a coreflection of C along $\underline{\mathcal{C}}^- \hookrightarrow \underline{\mathcal{C}}$. Thus K_C is uniquely determined up to a canonical isomorphism in $\underline{\mathcal{C}}^-$. Moreover if we allow the axiom of choice, we obtain a right adjoint of $\underline{\mathcal{C}}^- \hookrightarrow \underline{\mathcal{C}}$.

Lemma 4.6. *In the notation of Construction 4.2, if C belongs to $\mathcal{C}^-$, then Z_C belongs to $\mathcal{H}$.*

Dually, $C \in \mathcal{C}^+$ implies $K_C \in \mathcal{H}$ (in the notation of Remark 4.5).

Proof. We only show the former half. The latter is shown dually. Let

$$\begin{array}{c} U[-1] \xrightarrow{u} C \xrightarrow{v} V \rightarrow U \\ U_Z[-1] \xrightarrow{u_Z} Z_C \xrightarrow{v_Z} V_Z \rightarrow U_Z \end{array}$$

be distinguished triangles satisfying $U, U_Z \in \mathcal{U}$ and $V, V_Z \in \mathcal{V}$. By assumption, $V \in \mathcal{W}$. It suffices to show $V_Z \in \mathcal{W}$.

In the notation of Construction 4.2, complete $z_C \circ u: U[-1] \rightarrow Z_C$ into a distinguished triangle

$$U[-1] \xrightarrow{z_C \circ u} Z_C \rightarrow Q \rightarrow U.$$

By the octahedron axiom, we also have a distinguished triangle

$$V \rightarrow Q \rightarrow U'_C \rightarrow V[1],$$

and thus $Q \in \mathcal{U}$.

Since $\mathcal{C}(U[-1], V_Z) = 0$, we obtain a morphism of triangles

$$\begin{array}{ccccccc} U[-1] & \longrightarrow & Z_C & \longrightarrow & Q & \longrightarrow & U \\ \downarrow & & \Downarrow & & \downarrow & & \downarrow \\ U_Z[-1] & \longrightarrow & Z_C & \xrightarrow{v_Z} & V_Z & \longrightarrow & U_Z \end{array}$$

Since $\mathcal{C}(Q, V_Z) = 0$ by Lemma 2.11, it follows $\underline{v_Z} = 0$. For any $V^\dagger \in \mathcal{V}$ and any $v^\dagger \in \mathcal{C}(V_Z, V^\dagger[1])$ we have $\underline{v^\dagger \circ v_Z} = \underline{v^\dagger} \circ 0 = 0$ in $\mathcal{C}(Z_C, V^\dagger[1])$. By Proposition 2.10, we obtain $v^\dagger \circ v_Z = 0$. Thus $v^\dagger$ factors through U_Z , which means $v^\dagger = 0$ since $\mathcal{C}(U_Z, V^\dagger[1]) = 0$.

Thus we have $\mathcal{C}(V_Z, \mathcal{V}[1]) = 0$, namely $V_Z \in \mathcal{W}$. □

5. EXISTENCE OF (CO-)KERNELS

Lemma 5.1. *Let $A \xrightarrow{f} B$ be any morphism in $\mathcal{C}$. Take a diagram*

$$\begin{array}{ccc} U_A[-1] & & \\ \downarrow u_A & \searrow \circlearrowleft & \\ A & \xrightarrow{f} & B \\ \downarrow v_A & & \searrow m_f \\ V_A & & M_f \end{array}$$

where

$$\begin{aligned} U_A[-1] &\xrightarrow{u_A} A \xrightarrow{v_A} V_A \rightarrow U_A \\ U_A[-1] &\rightarrow B \xrightarrow{m_f} M_f \rightarrow U_A \end{aligned}$$

are distinguished triangles, satisfying $U_A \in \mathcal{U}_A, V_A \in \mathcal{V}$. Then we have the following.

- (1) $A \in \mathcal{C}^- \implies \underline{m_f} \circ \underline{f} = 0$,
- (2) $B \in \mathcal{C}^- \implies \overline{M_f} \in \mathcal{C}^-$.

Proof. (1) This immediately follows from $V_A \in \mathcal{W}$.

(2) Take distinguished triangles

$$\begin{aligned} U_B[-1] &\rightarrow B \rightarrow V_B \rightarrow U_B \\ U_M[-1] &\rightarrow M_f \rightarrow V_M \rightarrow U_M \\ (U_B, U_M &\in \mathcal{U}, \quad V_B, V_M \in \mathcal{V}). \end{aligned}$$

By assumption, $V_B \in \mathcal{W}$. It suffices to show $V_M \in \mathcal{W}$. Since $\mathcal{C}(U_B[-1], V_M) = 0$, there exists a morphism of triangles

$$\begin{array}{ccccccc} U_B[-1] & \xrightarrow{u_B} & B & \xrightarrow{v_B} & V_B & \longrightarrow & U_B \\ m_U \downarrow & & \downarrow & \circlearrowleft & \downarrow & & \downarrow \\ U_M[-1] & \xrightarrow{u_M} & M_f & \xrightarrow{v_M} & V_M & \longrightarrow & U_M \end{array}$$

Let $V^\dagger$ be any object in $\mathcal{V}$, and $v^\dagger \in \mathcal{C}(V_M, V^\dagger[1])$ be any morphism. It suffices to show $v^\dagger = 0$. Since $v^\dagger \circ m_V \in \mathcal{C}(V_B, V^\dagger[1]) = 0$, we have $v^\dagger \circ v_M \circ m_f = 0$.

$$\begin{array}{ccccc} B & \xrightarrow{v_B} & V_B \in \mathcal{W} & & \\ m_f \downarrow & & \downarrow m_V & & \\ M_f & \xrightarrow{v_M} & V_M & \longrightarrow & U_M \\ \downarrow & & \downarrow v^\dagger & \searrow \circlearrowleft & \swarrow 0 \\ U_A & \xrightarrow{0} & V^\dagger[1] & & \end{array}$$

Thus $v^\dagger \circ v_M$ factors through U_A , which means $v^\dagger \circ v_M = 0$, since $\mathcal{C}(U_A, V^\dagger[1]) = 0$. Thus $v^\dagger$ factors through U_M , and $v^\dagger = 0$ follows from $\mathcal{C}(U_M, V^\dagger[1]) = 0$. $\square$

Dually, we have the following:

Remark 5.2. For any morphism $A \xrightarrow{f} B$ in $\mathcal{C}$, consider a diagram

$$\begin{array}{ccccc}
 L_f & & & U_B & \\
 \searrow \ell_f & & & \downarrow u_B & \\
 & A & \xrightarrow{f} & B & \\
 & \searrow v_B \circ f & & \downarrow v_B & \\
 & & & V_B[1] &
 \end{array}$$

where

$$\begin{aligned}
 U_B &\xrightarrow{u_B} B \xrightarrow{v_B} V_B[1] \rightarrow U_B[1] \\
 V_B &\rightarrow L_f \xrightarrow{\ell_f} A \rightarrow V_B[1]
 \end{aligned}$$

are distinguished triangles satisfying $U_B \in \mathcal{U}, V_B \in \mathcal{V}$. Then, $A \in \mathcal{C}^+$ implies $L_f \in \mathcal{C}^+$, and $B \in \mathcal{C}^+$ implies $\underline{f} \circ \underline{\ell}_f = 0$.

Proposition 5.3. *Let $A \xrightarrow{f} B$ be any morphism in $\mathcal{C}$. Then $m_f: B \rightarrow M_f$ in Lemma 5.1 satisfies the following property:*

(*) *For any $C \in \mathcal{C}$ and any morphism $g \in \mathcal{C}(B, C)$ satisfying $\underline{g} \circ \underline{f} = 0$, there exists a morphism $c \in \mathcal{C}(M_f, C)$ such that $c \circ m_f = g$.*

$$\begin{array}{ccccc}
 A & \xrightarrow{f} & B & \xrightarrow{g} & C \\
 & & \searrow m_f & \nearrow c & \\
 & & M_f & &
 \end{array}$$

Moreover if $C \in \mathcal{C}^+$, then $\underline{c} \in \underline{\mathcal{C}}(M_f, C)$ satisfying

$$\underline{c} \circ \underline{m}_f = \underline{g}$$

is unique in $\underline{\mathcal{C}}(M_f, C)$. The dual statement also holds for L_f in Remark 5.2.

Proof. First we show the existence. By Proposition 2.10, $\underline{g} \circ \underline{f} \circ \underline{u}_A = 0$ means $g \circ f \circ u_A = 0$. Thus there exists $c \in \mathcal{C}(M_f, C)$ such that $c \circ m_f = g$.

$$\begin{array}{ccccc}
 U_A[-1] & & & & \\
 \downarrow u_A & \searrow f \circ u_A & & & \\
 & A & \xrightarrow{f} & B & \\
 & & & \searrow m_f & \\
 & & & M_f &
 \end{array}$$

To show the uniqueness, let

$$U_C \xrightarrow{u_C} C \xrightarrow{v_C} V_C[1] \rightarrow U_C[1]$$

be a distinguished triangle with $U_C \in \mathcal{W}, V_C \in \mathcal{V}$. Suppose $c, c' \in \mathcal{C}(M_f, C)$ satisfy

$$\underline{c} \circ \underline{m}_f = \underline{g} = \underline{c'} \circ \underline{m}_f.$$

By Proposition 2.10, $\underline{v}_C \circ (\underline{c} - \underline{c}') \circ \underline{m}_f = 0$ means $v_C \circ (c - c') \circ m_f = 0$. Thus $v_C \circ (c - c')$ factors through U_A , and thus $v_C \circ (c - c') = 0$ follows from $\mathcal{C}(U_A, V_C[1]) = 0$. Thus $c - c'$ factors through $U_C \in \mathcal{W}$, which means $\underline{c} = \underline{c}'$.

$$\begin{array}{ccccc} B & \xrightarrow{m_f} & M_f & \longrightarrow & U_A \\ \downarrow & \circlearrowleft & \downarrow c-c' & \circlearrowleft & \downarrow 0 \\ U_C & \xrightarrow{u_C} & C & \xrightarrow{v_C} & V_C[1] \end{array}$$

□

Corollary 5.4. *In $\underline{\mathcal{H}}$, any morphism has a cokernel and a kernel.*

Proof. We only show the construction of the cokernel. For any $A, B \in \mathcal{H}$ and any $f \in \mathcal{C}(A, B)$, define $m_f: B \rightarrow M_f$ as in Lemma 5.1. Since $A, B \in \mathcal{C}^-$, it follows

$$\underline{m}_f \circ \underline{f} = 0, \quad M_f \in \mathcal{C}^-$$

by Lemma 5.1. By Proposition 4.3, there exists $z_M: M_f \rightarrow Z_M$ which gives a reflection $\underline{z}_M: M_f \rightarrow Z_M$ of M_f along $\underline{\mathcal{C}}^+ \hookrightarrow \underline{\mathcal{C}}$. By Lemma 4.6, Z_M satisfies $Z_M \in \mathcal{H}$.

We claim that $\underline{z}_M \circ \underline{m}_f: B \rightarrow Z_M$ is the cokernel of $\underline{f}$. Let S be any object in $\mathcal{H}$, and let $s \in \mathcal{C}(B, S)$ be any morphism satisfying $\underline{s} \circ \underline{f} = 0$.

$$\begin{array}{c} \begin{array}{ccc} & \xrightarrow{0} & S \\ & \circlearrowleft & \uparrow \underline{s} \\ A & \xrightarrow{\underline{f}} & B \\ & \searrow \underline{m}_f & \\ & M_f & \\ & \searrow \underline{z}_M & \\ & Z_M & \end{array} \end{array}$$

It suffices to show that there uniquely exists $\underline{t} \in \underline{\mathcal{H}}(Z_M, S)$ such that

$$\underline{t} \circ \underline{z}_M \circ \underline{m}_f = \underline{s}.$$

This follows immediately from Proposition 4.3 and Proposition 5.3. □

6. ABELIANESS OF THE HEART

In this section, as the main theorem, we show that the heart $\underline{\mathcal{H}}$ becomes an abelian category, for any cotorsion pair $(\mathcal{U}, \mathcal{V})$. Although propositions and lemmas in this section could be applied for objects in $\mathcal{C}^+$ or $\mathcal{C}^-$ (with certain modifications of the statement), we mainly consider objects in $\mathcal{H}$.

Proposition 6.1. *Let $B, C \in \mathcal{H}$, and let*

$$A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} A[1]$$

be a distinguished triangle in $\mathcal{C}$. Let $m_g: C \rightarrow M_g$ be as in Lemma 5.1. Then the following are equivalent.

- (1) $\underline{g}$ is epimorphic in $\underline{\mathcal{H}}$.
- (2) $\underline{g}$ is epimorphic in $\underline{\mathcal{C}}^+$.
- (3) M_g satisfies $\underline{\mathcal{C}}(M_g, \underline{\mathcal{C}}^+) = 0$.
- (4) M_g satisfies $\mathcal{C}(M_g, \mathcal{V}[1]) = 0$.

The dual statement also holds for monomorphisms.

Proof. First, we show the equivalence of (1) and (2). Obviously, (2) implies (1). To show the converse, let S be any object in $\mathcal{C}^+$, and $\underline{s} \in \underline{\mathcal{C}}(C, S)$ be any morphism. Let $k_S: K_S \rightarrow S$ be the morphism defined in Remark 4.5, which gives a coreflection of S along $\underline{\mathcal{C}}^- \hookrightarrow \underline{\mathcal{C}}$. By Lemma 4.6, $K_S \in \mathcal{H}$.

By Remark 4.5, there exists $\underline{j} \in \underline{\mathcal{C}}^-(C, K_S)$ such that $\underline{k}_S \circ \underline{j} = \underline{s}$,

$$\begin{array}{ccccc} & & K_S & & \\ & \nearrow \underline{j} & \circlearrowleft & \searrow \underline{k}_S & \\ B & \xrightarrow{\underline{g}} & C & \xrightarrow{\underline{s}} & S \end{array}$$

and we have

$$\begin{aligned} \underline{j} = 0 &\iff \underline{k}_S \circ \underline{j} = 0 \\ \underline{j} \circ \underline{g} = 0 &\iff \underline{k}_S \circ (\underline{j} \circ \underline{g}) = 0. \end{aligned}$$

By (1), we have

$$\underline{j} = 0 \iff \underline{j} \circ \underline{g} = 0.$$

Thus $\underline{s} = 0$ if and only if $\underline{s} \circ \underline{g} = 0$, i.e., $\underline{g}$ is epimorphic in $\underline{\mathcal{C}}^+$.

Second, we show that (2) implies (3). By Proposition 5.3, for any $S \in \underline{\mathcal{C}}^+$ we have an isomorphism

$$-\circ \underline{m}_g: \underline{\mathcal{C}}(M_g, S) \xrightarrow{\cong} \{\underline{s} \in \underline{\mathcal{C}}(C, S) \mid \underline{s} \circ \underline{g} = 0\}.$$

Third, we show that (3) implies (4). This immediately follows from Proposition 2.10, since $\mathcal{V}[1] \subseteq \mathcal{C}^+$.

Finally, we show that (4) implies (3). Suppose $\mathcal{C}(M_g, \mathcal{V}[1]) = 0$. For any $S \in \mathcal{C}^+$, let

$$U_S \xrightarrow{u_S} S \xrightarrow{v_S} V_S[1] \rightarrow U_S[1]$$

be a distinguished triangle satisfying $U_S \in \mathcal{W}, V_S \in \mathcal{V}$. By $\mathcal{C}(M_g, \mathcal{V}[1]) = 0$, any morphism $s \in \mathcal{C}(M_g, S)$ factors through $U_S \in \mathcal{W}$, which means $\underline{s} = 0$. $\square$

Lemma 6.2. Let $B, C \in \mathcal{H}$, and let

$$A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} A[1]$$

be a distinguished triangle in $\mathcal{C}$. Take a diagram

$$\begin{array}{ccccc} U_A[-1] & \xrightarrow{f_U} & U_B[-1] & & \\ \downarrow u_A & \circlearrowleft & \downarrow u_B & & \\ A & \xrightarrow{f} & B & \xrightarrow{g} & C \\ \downarrow v_A & \circlearrowleft & \downarrow v_B & & \\ V_A & \xrightarrow{f_V} & V_B & & \end{array}$$

where

$$\begin{aligned} U_A[-1] &\xrightarrow{u_A} A \xrightarrow{v_A} V_A \rightarrow U_A \\ U_B[-1] &\xrightarrow{u_B} B \xrightarrow{v_B} V_B \rightarrow U_B \end{aligned}$$

are distinguished triangles satisfying $U_A, U_B \in \mathcal{U}, V_A \in \mathcal{V}$ and $V_B \in \mathcal{W}$.

If $\underline{g}$ is an epimorphism in $\underline{\mathcal{H}}$, then there exists a morphism $s \in \mathcal{C}(V_B, V_A)$ such that

$$s \circ v_B \circ f = v_A.$$

Proof. Let $C \xrightarrow{m_g} M_g$ be as in Lemma 5.1. Remark h factors m_g .

$$\begin{array}{ccc} C & \xrightarrow{h} & A[1] \\ & \searrow m_g & \swarrow \exists \\ & M_g & \end{array}$$

By Proposition 6.1, $\mathcal{C}(M_g[-1], V_A) \cong \mathcal{C}(M_g, V_A[1]) = 0$, and thus $v_A \circ h[-1] = 0$.

$$\begin{array}{ccc} C[-1] & \xrightarrow{h[-1]} & A \\ m_g[-1] \downarrow & \swarrow \circlearrowleft & \downarrow v_A \\ M_g[-1] & \xrightarrow{0} & V_A \end{array}$$

Thus there exists $q \in \mathcal{C}(B, V_A)$ such that $q \circ f = v_A$.

$$\begin{array}{ccccc} & & & & U_B[-1] \\ & & & & \downarrow u_B \\ C[-1] & \xrightarrow{h[-1]} & A & \xrightarrow{f} & B \\ & \searrow \circlearrowleft & \downarrow v_A & \swarrow \circlearrowleft & \downarrow v_B \\ & 0 & V_A & \xrightarrow{f_V} & V_B \\ & & & & \uparrow q \end{array}$$

Since $\mathcal{C}(U_B[-1], V_A) = 0$, we have $q \circ u_B = 0$, and thus there exists $s \in \mathcal{C}(V_B, V_A)$ such that $s \circ v_B = q$. Thus we obtain $s \circ v_B \circ f = q \circ f = v_A$. $\square$

Lemma 6.3. *In the notation of Lemma 6.2, if $\underline{g}$ is epimorphic in $\underline{\mathcal{H}}$, then A belongs to $\mathcal{C}^-$.*

Proof. We use the notation of Lemma 6.2. By the same lemma, there exists $s \in \mathcal{C}(V_B, V_A)$ such that $s \circ v_B \circ f = v_A$.

It suffices to show $V_A \in \mathcal{W}$. By assumption $V_B \in \mathcal{W}$ and thus for any $V^\dagger \in \mathcal{V}$ and any $v^\dagger \in \mathcal{C}(V_A, V^\dagger[1])$, we have $v^\dagger \circ s = 0$. Thus $v^\dagger \circ v_A = v^\dagger \circ s \circ v_B \circ f = 0$, and $v^\dagger$ factors through U_A , which implies $v^\dagger = 0$.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ v_A \downarrow & \circlearrowleft & \downarrow v_B \\ V_A & \xleftarrow{s} & V_B \\ & \searrow v^\dagger & \swarrow 0 \\ & \circlearrowleft & \\ U_A & \xrightarrow{0} & V^\dagger[1] \end{array}$$

$\square$

Theorem 6.4. *For any cotorsion pair $(\mathcal{U}, \mathcal{V})$ in $\mathcal{C}$, its heart*

$$\underline{\mathcal{H}} = (\mathcal{C}^+ \cap \mathcal{C}^-) / (\mathcal{U} \cap \mathcal{V})$$

is an abelian category.

Proof. Since $\underline{\mathcal{H}}$ is an additive category with (co-)kernels (Corollary 5.4), it remains to show the following:

- (1) If $\underline{g}$ is epimorphic in $\underline{\mathcal{H}}$, then $\underline{g}$ is a cokernel of some morphism in $\underline{\mathcal{H}}$.
- (2) If $\underline{g}$ is monomorphic in $\underline{\mathcal{H}}$, then $\underline{g}$ is a kernel of some morphism in $\underline{\mathcal{H}}$.

Since (2) can be shown dually, we only show (1).

Complete g into a distinguished triangle

$$A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} A[1]$$

in $\mathcal{C}$. By Lemma 6.3, we have $A \in \mathcal{C}^-$. By Proposition 4.3, there exists

$$z_A: A \rightarrow Z_A$$

which gives a reflection $\underline{z}_A: A \rightarrow Z_A$ of A along $\underline{\mathcal{C}}^+ \hookrightarrow \underline{\mathcal{C}}$. Moreover by Lemma 4.6, Z_A belongs to $\mathcal{H}$. Since B belongs to $\mathcal{C}^+$, there exists $b \in \mathcal{C}(Z_A, B)$ such that $b \circ z_A = f$.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ & \searrow z_A & \nearrow b \\ & Z_A & \end{array}$$

We claim that $\underline{g} = \text{cok}(\underline{b})$.

Let S be any object in $\mathcal{H}$, and $s \in \mathcal{C}(B, S)$ be any morphism. By Proposition 4.3, $\underline{s} \circ \underline{f} = 0$ if and only if $\underline{s} \circ \underline{b} = 0$. In particular, $\underline{g} \circ \underline{b} = 0$.

So it suffices to show for any s satisfying $\underline{s} \circ \underline{f} = 0$, there uniquely exists $\underline{c} \in \underline{\mathcal{C}}(C, S)$ such that $\underline{c} \circ \underline{g} = \underline{s}$.

Uniqueness immediately follows from the fact that $\underline{g}$ is epimorphic. So it remains to show the existence of $\underline{c}$. Since $A \in \mathcal{C}^-$, there exists a distinguished triangle

$$U_A[-1] \xrightarrow{u_A} A \xrightarrow{v_A} V_A \rightarrow U_A$$

with $U_A \in \mathcal{U}, V_A \in \mathcal{W}$. By assumption we have $\underline{s} \circ \underline{f} \circ \underline{u}_A = 0$, and this means $s \circ f \circ u_A = 0$ by Proposition 2.10. So $s \circ f$ factors through V_A , and we obtain a morphism of triangles

$$\begin{array}{ccccccc} A & \xrightarrow{f} & B & \xrightarrow{g} & C & \xrightarrow{h} & A[1] \\ v_A \downarrow & \circlearrowleft & \downarrow s & \circlearrowleft & \downarrow t & \circlearrowleft & \downarrow -v_A[1] \\ V_A & \xrightarrow{i} & S & \xrightarrow{j} & T & \longrightarrow & V_A[1] \end{array}$$

where $V_A \xrightarrow{i} S \xrightarrow{j} T \rightarrow V_A[1]$ is a distinguished triangle. By the same argument as in the proof of Lemma 6.2 (since $\mathcal{C}(M_g, V_A[1]) = 0$), we have $v_A[1] \circ h = 0$. Thus t factors through S , namely there exists $c \in \mathcal{C}(C, S)$ such that $j \circ c = t$. Since

$j \circ (s - c \circ g) = j \circ s - t \circ g = 0$, there exists $s' \in \mathcal{C}(B, V_A)$ such that

$$i \circ s' = s - c \circ g.$$

$$\begin{array}{ccccc} & & B & \xrightarrow{g} & C \\ & \swarrow s' & \downarrow s & \searrow c & \downarrow t \\ V_A & \xrightarrow{i} & S & \xrightarrow{j} & T \end{array}$$

Since $V_A \in \mathcal{W}$, this means $\underline{s} = \underline{c} \circ \underline{g}$. $\square$

7. EXISTENCE OF ENOUGH PROJECTIVES/INJECTIVES

Lemma 7.1. *For any cotorsion pair $(\mathcal{U}, \mathcal{V})$, the following are equivalent.*

- (1) $\mathcal{U} \subseteq \mathcal{V}$.
- (2) $\mathcal{W} = \mathcal{U}$.
- (3) $\mathcal{C}^+ = \mathcal{C}$.

Proof. Left to the reader. $\square$

Corollary 7.2. *If a cotorsion pair $(\mathcal{U}, \mathcal{V})$ satisfies $\mathcal{U} \subseteq \mathcal{V}$, then we have $\mathcal{U}[-1] \subseteq \mathcal{H}$.*

Proof. This immediately follows from $\mathcal{U}[-1] \subseteq \mathcal{C}^-$ (Remark 3.6) and $\mathcal{C} = \mathcal{C}^+$ (Lemma 7.1). $\square$

Proposition 7.3. *Let $(\mathcal{U}, \mathcal{V})$ be a cotorsion pair satisfying $\mathcal{U} \subseteq \mathcal{V}$. If an object $P \in \mathcal{H}$ lies in the image of $\mathcal{U}[-1]$ (i.e. $P \in \mathcal{U}[-1]/\mathcal{W}$), then P is projective in $\underline{\mathcal{H}}$.*

Proof. Let B and C be any objects in $\mathcal{H}$ and let $p \in \mathcal{C}(P, C)$ be any morphism.

Let $g \in \mathcal{C}(B, C)$ be any morphism which is epimorphic in $\underline{\mathcal{H}}$, and take a distinguished triangle

$$A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} A[1].$$

By Proposition 6.1 and Lemma 7.1, $\underline{g}$ is epimorphic in $\underline{\mathcal{C}}$. By Fact 2.9, this is equivalent to $\underline{h} = 0$. Thus h factors through some $W \in \mathcal{W}$. Since $\mathcal{C}(P, W) \subseteq \text{Ext}^1(\mathcal{U}, \mathcal{V}) = 0$, we have $h \circ p = 0$, and p factors through B as desired. $\square$

Corollary 7.4. *If a cotorsion pair $(\mathcal{U}, \mathcal{V})$ satisfies $\mathcal{U} \subseteq \mathcal{V}$, then its heart $\underline{\mathcal{H}}$ has enough projectives.*

Proof. By definition, for any $C \in \mathcal{H}$, there exists a distinguished triangle

$$U[-1] \xrightarrow{u} C \xrightarrow{v} V \rightarrow U$$

with $U \in \mathcal{U}, V \in \mathcal{V}$. Since $\underline{v} = 0$ by $\underline{\mathcal{C}}(C, V) = 0$ (Lemma 2.11), $\underline{u}$ is epimorphic in $\underline{\mathcal{C}}$, and thus in $\underline{\mathcal{H}}$. Thus Corollary 7.4 follows from Corollary 7.2 and Proposition 7.3. $\square$

Dually, we have the following.

Corollary 7.5. *If a cotorsion pair $(\mathcal{U}, \mathcal{V})$ satisfies $\mathcal{V} \subseteq \mathcal{U}$, then its heart $\underline{\mathcal{H}}$ has enough injectives.*

8. EXAMPLE ARISING FROM AN n -CLUSTER TILTING SUBCATEGORY

Definition 8.1. (§§5.1 in [KR]) A full additive thick subcategory $\mathcal{T} \subseteq \mathcal{C}$ is an n -cluster tilting subcategory if it satisfies the following.

- (1) $\mathcal{T}$ is functorially finite.
- (2) An object $C \in \mathcal{C}$ belongs to $\mathcal{T}$ if and only if $\text{Ext}^\ell(C, \mathcal{T}) = 0$ ($0 < \forall \ell < n$)
- (3) An object $C \in \mathcal{C}$ belongs to $\mathcal{T}$ if and only if $\text{Ext}^\ell(\mathcal{T}, C) = 0$ ($0 < \forall \ell < n$)

Definition 8.2. Let $\mathcal{T}$ be an $n+1$ -cluster tilting subcategory. For any pair of integers i and j satisfying $i \leq j$, put

$$\begin{aligned} \mathcal{T}_{[i,j]} &:= \{C \in \mathcal{C} \mid \text{Ext}^\ell(C, \mathcal{T}) = 0 \ (j+1 \leq \forall \ell \leq n)\}, \\ \mathcal{T}^{[i,j]} &:= \{C \in \mathcal{C} \mid \text{Ext}^\ell(\mathcal{T}, C) = 0 \ (-j \leq \forall \ell \leq -i)\}. \end{aligned}$$

In particular, if $j-i \geq n$, then $\mathcal{T}_{[i,j]} = \mathcal{T}^{[i,j]} = \mathcal{C}$.

Example 8.3. Let $\mathcal{T}$ be an $n+1$ -cluster tilting subcategory. For any integer ϖ satisfying $0 \leq \varpi \leq n-1$, put

$$\begin{aligned} \mathcal{T}_\varpi &:= \mathcal{T}_{[0,\varpi]}, \\ \mathcal{T}^\varpi &:= \mathcal{T}^{[\varpi, n-1]}. \end{aligned}$$

Then it can be shown that $(\mathcal{T}_\varpi, \mathcal{T}^\varpi)$ is a cotorsion pair (Theorem 3.1 in [IY]). We can calculate its heart $\underline{\mathcal{H}}_\varpi$ as follows.

$$\begin{aligned} \mathcal{C}^+ &= \mathcal{T}_{[\varpi, n]}, \quad \mathcal{C}^- = \mathcal{T}^{[-1, \varpi]}, \quad \mathcal{W} = \mathcal{T}[\varpi], \\ \underline{\mathcal{H}}_\varpi &= (\mathcal{T}_{[\varpi, n]} \cap \mathcal{T}^{[-1, \varpi]}) / \mathcal{T}[\varpi]. \end{aligned}$$

Thus $(\mathcal{T}_{[\varpi, n]} \cap \mathcal{T}^{[-1, \varpi]}) / \mathcal{T}[\varpi]$ is an abelian category for each $\varpi \in [0, n-1]$, as shown in Corollary 6.4 in [IY]. If $n=1$ and $\varpi=0$, this is nothing other than the case of a cluster tilting subcategory.

Moreover in Corollary 6.4 in [IY], it was shown that these $(\mathcal{T}_{[\varpi, n]} \cap \mathcal{T}^{[-1, \varpi]}) / \mathcal{T}[\varpi]$ are mutually equivalent for $\varpi \in [0, n-1]$. We abbreviate these equivalent abelian categories by $\underline{\mathcal{H}}$.

Since $(\mathcal{T}_0, \mathcal{T}^0)$ satisfies $\mathcal{T}_0 = \mathcal{T} \subseteq \mathcal{T}^0$, we see that $\underline{\mathcal{H}}$ has enough projectives by Corollary 7.4. Dually, since $(\mathcal{T}_{n-1}, \mathcal{T}^{n-1})$ satisfies $\mathcal{T}^{n-1} = \mathcal{T}[n-1] \subseteq \mathcal{T}_{n-1}$, we see $\underline{\mathcal{H}}$ has enough injectives by Corollary 7.5.

REFERENCES

- [BBD] Beilinson, A. A.; Bernstein, J.; Deligne, P.: *Faisceaux pervers* (French) [Perverse sheaves] Analysis and topology on singular spaces, I (Luminy, 1981), 5–171, Astérisque, **100**, Soc. Math. France, Paris, 1982.
- [BR] Beligiannis, A.; Reiten, I.: *Homological and homotopical aspects of torsion theories* (English summary), Mem. Amer. Math. Soc. **188** (2007), no. 883, viii+207 pp.
- [B] Borceux, F.: *Handbook of categorical algebra 1, Basic category theory*, Encyclopedia of Mathematics and its Applications, **50**. Cambridge University Press, Cambridge, 1994. xvi+345 pp.
- [BMR] Buan, A. B.; Marsh, R. J.; Reiten, I.: *Cluster-tilted algebras*, Trans. Amer. Math. Soc. **359** (2007), no. 1, 323–332.
- [IY] Iyama, O.; Yoshino, Y.: *Mutation in triangulated categories and rigid Cohen-Macaulay modules* (English summary), Invent. Math. **172** (2008), no. 1, 117–168.
- [KR] Keller, B.; Reiten, I.: *Cluster-tilted algebras are Gorenstein and stably Calabi-Yau*, Adv. Math. **211** (2007), no. 1, 123–151.
- [KZ] Koenig, S.; Zhu, B.: *From triangulated categories to abelian categories: cluster tilting in a general framework* (English summary), Math. Z. **258** (2008), no. 1, 143–160.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO 3-8-1 KOMABA,
MEGURO, TOKYO, 153-8914 JAPAN
E-mail address: `deutsche@ms.u-tokyo.ac.jp`